



MINIMIZING DISEASES OF PEANUT IN THE SOUTHEASTERN UNITED STATES

University of Georgia

Robert Kemerait, Albert Culbreath, Eric Prostko, Tim Brenneman,
Scott Tubbs, Rajagopalbabu Srinivasan, Mark Abney,
Scott Monfort, Cristiane Pilon, and Sudeep Bag

University of Florida

Barry Tillman, Nicholas Dufault, Michael Mulvaney and Ian Small

Auburn University

Austin Hagan, Charles Chen, Alana Jacobson, Kris Balkcom,
and Amanda Strayer-Scherer

Clemson University

Dan Anco

Mississippi State University

Brendan Zurweller

ASSESS DISEASE RISK IN YOUR FIELD AND DEVELOP A PEANUT RX


This worksheet will lead you through the four-step process of determining your disease risk level in order to customize a Peanut Rx[™] for your individual field using the reverse side of this worksheet and with the assistance of your Syngenta representative.

For each of the risk index factors, identify which option best describes the situation for your field and add the index value associated with each choice to obtain your overall disease risk value. This worksheet does not contain all of the varieties included in the 2021 Peanut Rx or the notes that accompany each factor. To view the complete 2021 Peanut Rx, visit the University of Georgia peanut website at www.uga-peanutteam.com.

Assess Your Disease Risk

Variety Selection			
Variety ¹	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points White Mold
AU NPL 17 ²	10	15	15
Bailey ³	10	25	10
Florida Fancy ²	25	20	20
FloRun [™] 331 ^{1,2}	15	20	15
Georgia-06G	10	20	20
Georgia-07W	10	20	15
Georgia-09B ²	20	25	25
Georgia-12Y ²	5	15	10
Georgia-14N ^{2,4}	5	15	15
Georgia-16HQ ²	10	25	20
Georgia-18RU ¹	10	25	20
Georgia Green	30	20	25
Sullivan ²	10	25	15
Tifguard ⁴	10	15	15
TifNV-HiOL ^{2,4}	5	15	15
TUFRunner [™] 287 ^{1,2}	10	25	20
TUFRunner [™] 511 ^{1,2}	20	30	15
Planting Date			
Peanuts are planted:	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points White Mold
Prior to May 1	30	0	10
May 1 to May 10	15	5	5
May 11 to May 25	5	10	0
May 26 to June 10	10	15	0
After June 10	15	15	0
Plant Population (final stand, not seeding rate)			
Plant stand:	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points White Mold
Less than 3 plants/ft	25	NA	0
3 to 4 plants/ft (3)	10 (15)	NA	0 (0)
More than 4 plants/ft	5	NA	5
At-plant Insecticide			
Insecticide used	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points White Mold
None	15	5	NA
Other than Thimet [®] 20G	15	5	NA
Velum Total	15	0	NA
Thimet 20G	5	0	NA
Row Pattern			
Peanuts are planted in:	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points White Mold
Single rows	10	0	5
Twin rows	5	0	0
Tillage			
Tillage type	Spotted Wilt Points	Leaf Spot Points	Soilborne Disease Points White Mold
Conventional	15	10	0
Reduced	5	0	5

This guide was funded by:



College of Agricultural & Environmental Sciences
UNIVERSITY OF GEORGIA

Annual Publication AP-118



Calculate Your Severity Points

Fill in the following table to calculate your severity points for each of the four major peanut diseases given the 10 determining factors. Total each column to establish your disease index values.

Variety	Spotted Wilt	Leaf Spot	White Mold	Rhizoctonia Limb Rot
Planting Date				
Plant Population				
At-plant Insecticide				
Row Pattern				
Tillage				
Classic Herbicide				
Crop Rotation				
Field History				
Irrigation				
Total Index Value				

Interpret Your Risk Total

Point total range for tomato spotted wilt = 35-155.
Point total range for leaf spot = 10-105.
Point total range for white mold = 10-95.
Point total range for Rhizoctonia limb rot = 15-75.

High Risk	Spotted Wilt Points	Leaf Spot Points	Soilborne Points	
			White Mold	Limb Rot
High Risk for fungal diseases: Growers should always use full fungicide input program in a high-risk situation.	≥ 115	65-105	55-80	TBD
Moderate Risk	70-110	40-60	30-50	TBD
Medium Risk for fungal diseases: Growers can expect better performance from standard fungicide programs. Reduced fungicide programs in research studies have been successfully implemented when conditions are not favorable for disease spread.	≤ 65	10-35	10-25	TBD
Low Risk	≤ 65	10-35	10-25	TBD
Low Risk for fungal diseases: These fields are likely to have the least impact from fungal disease. Growers have made the management decisions which offer maximum benefit in reducing the potential for severe disease; these fields are strong candidates for modified disease management programs that require a reduced number of fungicide application.				

When tomato spotted wilt virus incidence is high statewide or in your region, even fields with a low risk level may experience significant losses. Consider the following recommendations to reduce your spotted wilt risk level:

- Use less susceptible varieties
- Adjust your planting date
- Consult the complete Peanut Rx for additional options that may also provide limited benefit

Develop Your Peanut Rx

Once you have calculated your total risk for each fungal disease, utilize the most conservative fungicide program as your guide for customizing a per-field prescription spray program.

Programs developed through the cooperation of

